ORIGINAL ARTICLE

Incidental poisoning of animals by carbamates in the Czech Republic

Ladislav Novotný¹, Jan Misík², Alena Honzlová⁴, Petr Ondráček⁵, Kamil Kuča^{1,3}, Oldřich Vávra⁴,
Václav Rachač⁶, Petr Chloupek⁵

¹Center of Advanced Studies, ²Department of Toxicology, Faculty of Military Health Sciences, University of Defence, Hradec Králové, Czech Republic

³University Hospital, Hradec Králové, Czech Republic

⁴State Veterinary Institute, Jihlava, Czech Republic

⁵Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic

⁶State Veterinary Administration of the Czech Republic, České Budějovice, Czech Republic

Received 16th September 2010.

Revised 20th October 2010.

Published online 23rd November 2010.

Summary

Illegal poisoning of wildlife and domestic animals is a worldwide issue. The carbamates primarily used as pesticides are often misused for such a purpose. In this study, 181 birds, mammals and baits were analysed over the period 2004–09 for possible intoxication by carbamates. Intoxication by carbamate carbofuran was diagnosed in 89 cases, and in another 19 cases (nine Wild Boars and 10 Bisons) intoxication with another carbamate – methomyl – was proven. Incidental ingestion of the marten bait was the main cause of intoxication. Although the distribution of carbofuran was prohibited in 2007, no decline in the number of intoxicated animals in the following two years was detected. New cases of intoxication by carbofuran are anticipated in the future until all remaining stock is expended.

Key words: pesticide poisoning; organophosphorous inhibitors; forensic medicine; acetylcholinesterase

INTRODUCTION

Pesticide poisoning of non target organisms is a worldwide issue and many cases are described (Roberston et al. 1992, Hornfeldt and Murphy 1997,

Xavier et al. 2002, Berny and Gaillet 2008). Carbamate compounds are a very important group of cholinesterase inhibitors and intoxications caused by these compounds represent the majority of pesticide poisoning among animals. According to present knowledge, carbofuran (2,3-dihydro-2,2-dimethyl-7-benzofuranylmethylcarbamate) is the most frequently encountered carbamate as a cause of intoxication. Its LD₅₀ is ranging from 3–19 mg.kg⁻¹ of body weight in many animal species (Wang et al. 2007).

Toxicologically, carbamates belong to the group of acylating (carbamylation) acetylcholinesterase (AChE) inhibitors which inhibit AChE by carbamylation of the serine hydroxyl group in the

✉ Ladislav Novotný, Faculty of Military Health Sciences, University of Defence, Třebešská 1575, 500 01 Hradec Králové, Czech Republic

novotnyl@pmfhk.cz

420 723 289 823

enzyme active site. Moreover, this inhibition leads to the accumulation of acetylcholine in the gap junction causing hyperstimulation of cholinergic receptors. “Overdose” with acetylcholine is followed by overstimulation of the receptors and finally it ends with a cholinergic crisis followed by muscarine, nicotine and central nervous signs, i.e. miosis, hypersecretion of exocrine glands, bradycardia, tonic and clonic convulsions (O’Malley 1997).

Although carbamate pesticides are less toxic than organophosphorus pesticides, many cases of animal poisonings caused by these agents are reported with increasing frequency (Balcom 1983, Elliot et al. 1996, Mineau et al. 1999, Wobeser et al. 2004).

In this study, we summarize the results from the analysis of 89 cases of positively tested birds (White-tailed Eagle, Golden Eagle, Common Buzzard, Rough-legged Buzzard, Hooded Crow, Common Raven, Common Magpie, Grey Partridge) and mammals (dog, Red Fox, Otter, Wild Boar, European Polecat, Marten) during the period 2004–09 poisoned by carbofuran and 19 cases of animals (Wild Boar, Bison) intoxicated with other carbamate methomyl (methyl *N* -{[(methylamino)carbonyl]oxy}ethanimidothioate).

MATERIAL AND METHODS

In this retrospective study, 181 cases of suspected intoxications from January 2004 to December 2009 were analysed. In these 181 cases, 59.0% were tissue samples (mostly liver), 31.0% were gastric content and 9.8% were baits (mostly eggs, meat and meat products). Baits had been laid in forests and parks, or thrown into private gardens.

White-tailed Eagle, Golden Eagle, Common Buzzard, Rough-legged Buzzard, Hooded Crow, Common Raven, Common Magpie, Grey Partridge, dog, Red Fox, Otter, Wild Boar, European Polecat, Marten, Bison were among the intoxicated animals.

Necropsy and histopathology were performed on entire animals (six dogs and three otters) delivered with a no longer *post mortem* interval than 24 hours.

The analysis was conducted with the available chromatographic techniques at the State Veterinary Institute in Jihlava Czech Republic, using tissue samples (liver), solid (bait, undefined material) or liquid samples (gastric content). Samples were extracted by petroleum ether and after evaporation on the rotary vacuum evaporator and dissolution in isoctane were analysed by the gas chromatography method with a nitrogen-phosphorous detector (Lee and Westcott 1983).

RESULTS

The numbers of examined samples and positive cases in each year of the study are given in Fig. 1. The presence of carbofuran could be verified in a total of 89 samples. This means that there were approximately half (49.2%) of the samples positive out of 181 in 6 years. In 2004 only three samples were examined. Two of these three samples originally from birds of prey were positive for carbofuran – a White-tailed Eagle (*Haliaeetus albicilla*) and a Rough-legged Buzzard (*Buteo lagopus*). In 2005, the percentage of positive samples was 45.5% (10/25), as in 2007 and 2008 [2007 44.0% (11/25), 2008 46.2% (18/39)]. In 2006, the percentage of positive samples was somewhat higher [67.9% (36/53)], whereas some decline was obvious in 2009 [30.8% (12/39)]. Birds of prey were the most common victims of carbofuran (Fig. 2) – 22 Buzzards (*Buteo* sp.) and 8 endangered Eagles (*Haliaeetus* sp., *Aquila* sp.) were illegally poisoned during the studied time frame. Also dogs (n=12), Otters (n=10, *Lutra lutra*) and Corvids [n=10; Common Raven (*Corvus corax*), Common Magpie (*Pica pica*) and Hooded Crow (*Corvus cornix*)] were frequently involved. Other species, such as small carnivores [Marten (*Martes* sp.), European Polecat (*Mustela putorius*), Stoat (*Mustela erminea*)] or birds [Grey Partridge (*Perdix perdix*)] rarely show signs of intoxication by carbamates.

In 2007 and 2008, 19 samples were positively tested for another carbamate – methomyl (methyl *N* -{[(methylamino)carbonyl]oxy}ethanimidothioate). Nine samples were taken from Wild Boars (*Sus scrofa*), another ten samples were collected from Bison (*Bison* sp.) from a private farm in south Bohemia. Up to November 2008 a total of a 37 Bison died supposedly poisoned by methomyl (not all animals were investigated).

Necropsy revealed dried saliva around the oral cavity, congestion of the organs, and haemorrhagic necrosis of the small gut. Histopathology showed congestion of the kidney, liver and lung, granular dystrophy of liver and haemorrhagic necrosis of the small intestine.

DISCUSSION

Carbamate intoxication is the most common form of pesticide poisoning of animals in Europe and the US (Berny 2007, Wang et al. 2007). The most common route of exposure to pesticides is an ingestion of poisoned insects, carcasses, or grains intentionally treated with pesticides for bait (Reece and Handson

Fig.1. No. of samples analysed in the State Veterinary Institute Jihlava tested for carbofuran (black columns) and number of positive cases (gray columns).

Fig.2. Distribution of carbofuran intoxicated samples. Birds of prey were the victims of carbofuran in most cases.

1982). Organophosphorous and carbamate pesticides present also a potential threat for humans mainly because of their wide use in agriculture. Thousands of intoxication cases in humans are recorded every year especially in developing countries (Jayaratnam 1990).

According to many studies, birds of prey are common victims of carbofuran poisoning (Balcom 1983, Elliot et al. 1996, Wobeser et al. 2004) and this was confirmed in our study. This could be caused by secondary poisoning when the raptor consumes a primary victim with unassimilated remains of pesticide in its alimentary tract (Dietrich et al. 1995, Hill 1995, Mineau et al. 1999, Berny 2007), but it is also possible that some of the baits are directed at birds of prey.

Compared to the high number of intoxicated birds we found only sporadic cases of small carnivore intoxication (except otters), although martens or foxes are thought to be the main target object of poisoners. This could be caused by the hidden behaviour of carnivores, so the real number of intoxications is expected to be much higher.

Also domesticated animals such as pets, mainly dogs, and livestock are at risk of being poisoned (Wang et al. 2007). The consumption of bait could be accidental during walks outdoors but there are also some cases of intentional poisoning (Novotný et al. 2003).

Although the use of carbofuran was prohibited by law in the Czech Republic on 13th December 2008 and its sale was stopped on 13th December 2007, new cases of intoxication by carbofuran are still expected in the future until all stocks are expended.

In organophosphorous and carbamate pesticide investigation, analysis of the brain and blood cholinesterase (ChE) activity is widely used as a diagnostic technique (Worek et al. 1999 Zdarova Karasova et al. 2009), but gas chromatography was used in this study. However, diagnostic interpretation of ChE data always requires knowledge of the normal value for each species (Blakley and Skelley 1988), and it may be confounded as well by changes in ChE levels caused by post mortem decomposition (Wobeser et al. 2004) and post mortem reactivation of carbamylated ChE (Hill 1989).

ACKNOWLEDGEMENT

The work was supported by the grant projects MSM0021620820 and OVUOFVZ200905.

REFERENCES

- Balcom R. Secondary poisoning of red-shouldered hawks with carbofuran. *J Wildl Manage.* 47: 1129–1132, 1983.
- Berny P. Pesticides and the intoxication of wild animals. *J Vet Pharmacol Ther.* 30: 93–100, 2007.
- Berny P, Gaillet JR. Acute poisoning of Red Kites (*Milvus milvus*) in France: Data from the SAGIR network. *J Wildl Dis.* 44: 417–426, 2008.
- Blakley BR, Skelley KW. Brain cholinesterase activity in animals and birds. *Vet Hum Toxicol.* 30: 329–331, 1988.
- Dietrich DR, Schmid P, Zweifel U, Schlatter C, Jenni-Eiermann S, Bachmann H, Bühler U, Zbinden N. Mortality of birds of prey following field application of granular carbofuran – a Case-study. *Arch Environ Contam Toxicol.* 29: 140–145, 1995.
- Elliot JE, Langelier KM, Mineau P, Wilson LK. Poisoning of bald eagles and red-tailed hawks by carbofuran and fensulfothion in the Fraser Delta of British Columbia, Canada. *J Wildl Dis.* 32: 486–491, 1996.
- Hill EF. Divergent effects of postmortem ambient temperature on organophosphorus- and carbamate-inhibited brain cholinesterase activity in birds. *Pestic Biochem Physiol.* 33: 264–275, 1989.
- Hill EF. Organophosphorous and carbamate pesticides. In Hoffman DJ, Rattner BA, Burton GH, Jr., Cairns J, Jr. (eds.): *Handbook of Ecotoxicology*, Lewis publ., Florida 1995, pp. 243–274.
- Hornfeldt CS, Murphy MJ. Poisonings of animals: the 1993–1994 report of the American Association of Poison Control Centers. *Vet Hum Toxicol.* 39: 361–365, 1997.
- Jayaratnam J. Pesticide poisoning as a global health problem. *World Health Stat Q.* 43: 139–144, 1990.
- Lee YW, Westcott ND. Direct analysis of carbofuran and its carbamate metabolites in rapeseed plants by nitrogen-phosphorus detector gas chromatography. *J Agric Food Chem.* 31: 92–96, 1983.
- Mineau P, Fletcher MR, Glazer LC, Thomas NJ, Brassard C, Wilson LK, Elliot JE, Lyon LA, Henny CJ, Bollinge T, Porter SL. Poisoning of raptors with organophosphorous pesticides with emphasis on Canada, US, and UK. *J Raptor Res.* 33: 1–37, 1999.
- Novotný L, Honzlová A, Ondráček P, Vávra O, Rachač V, Chloupek P. Intoxication of animals with carbofuran (in Czech). *Veterinářství* 53: 551–554, 2003.

- O'Malley M. Clinical evaluation of pesticide exposure and poisonings. *Lancet*. 349: 1161–1166, 1997.
- Reece RL, Handson P. Observations on the accidental poisoning of birds by organo-phosphate insecticides and other toxic substances. *Vet Rec*. 111: 453–455, 1982.
- Roberston ID, Leggoe M, Dorling PR, Shaw SE, Clark WT. A retrospective study of poisoning cases in dogs and cats: comparisons between a rural and an urban practice. *Aust Vet J*. 69: 194–195, 1992.
- Wang Y, Kruzik P, Helsberg A, Helsberg I, Rausch WD. Pesticide poisoning in domestic animals and livestock in Austria: a 6 years retrospective study. *Forensic Sci Int*. 169: 157–160, 2007.
- Wobeser G, Bollinger T, Leighton FA, Blakley B, Mineau P. Secondary poisoning of eagles following intentional poisoning of coyotes with anticholinesterase pesticides in western Canada. *J Wildl Dis*. 40: 163–172, 2004.
- Worek F, Mast U, Kiderlen D, Diepold C, Eyer P. Improved determination of acetylcholinesterase activity in human whole blood. *Clin Chim Acta*. 288: 73–90, 1999.
- Xavier FG, Kogika MM, Spinoia HS. Common causes of poisoning in dogs and cats in a Brazilian veterinary hospital 1998 to 2000. *Vet Hum Toxicol*. 44: 115–116, 2002.
- Zdarova Karasova J, Kassa J, Musilek K, Pohanka M, Novotny L, Kuca K. Effect of seven newly synthesized and currently available oxime cholinesterase reactivators on cyclosarin-intoxicated rats. *Int J Mol Sci*. 10: 3065–3075, 2009.