ORIGINAL ARTICLE

Docking studies and effects of *syn-anti* isomery of oximes derived from pyridine imidazol bicycled systems as potential human acetylcholinesterase reactivators

Ana Paula Guimarães^{1,2}, Tanos Celmar Costa França¹, Teodorico Castro Ramalho²,
Magdalena Nascimento Rennó³, Elaine Fontes Ferreira da Cunha², Karina Silvia Matos²,
Daiana Teixeira Mancini², Kamil Kuča^{4,5}

¹Laboratory of Molecular Modeling Applied to the Chemical and Biological Defense, Military Institute of Engineering, Rio de Janeiro, Brazil

²Chemistry Department, Federal University of Lavras, Lavras, Brazil

³Pharmacy Faculty, Federal University of Rio de Janeiro, Campus Macaé, Macaé, Brazil

⁴Department of Toxicology, Faculty of Military Health Sciences, University of Defence, Hradec Králové, Czech Republic

⁵University Hospital, Hradec Králové, Czech Republic

Received 15th November 2010.

Revised 10th February 2011.

Published online 7th April 2011.

Summary

In order to contribute to a better understanding of the mechanism of action of oximes, we evaluated the affinities of 10 new oximes, derived from pyridine-imidazol bicycled systems, for human acetylcholinesterase (*HssAChE*) complexed with tabun, by estimating their docking energy values and comparing of the values obtained to known oximes using the software Molegro Virtual Docker (MVD)[®]. We evaluated the influence of the position of the oxime group as substituent in the structures and, also, the influence of the oxime group *syn-anti* isomery on the docking score values for all the molecules studied. Results suggest that: the affinities of the 10 new oximes for the tabun inhibited *HssAChE* active site are better than pralidoxime's and similar to trimedoxime's; the *meta*-pralidoxime could have more affinity for the *HssAChE* active site and the oximes' *anti* isomers could present slightly better affinities for the *HssAChE* active site than the *syn* isomers.

Key words: acetylcholinesterase; docking studies; oximes; neurotoxic agents; theoretical calculation

INTRODUCTION

Despite the existence of many different oximes in use today against intoxication with neurotoxic agents, the

literature has not yet reported one able to act effectively against all the existing neurotoxic agents and oximes usually effective with one specific nerve agent can be completely ineffective with another. This probably happens because their mechanisms of action are not yet well elucidated (Ekström et al. 2007). In addition, other relevant factors: the adequate orientation of the phosphoryl bond inside the active site, the most suitable oxime charge, the most adequate angles for attacking the phosphorylated serine, the influence of the oxime's isomery, and the chemical environment of the oxime group, remain unknown despite the fact that they are recurrent issues in the literature (Bay et al. 1958, Smirnova et al.

✉ Teodorico Castro Ramalho, Chemistry Department, Federal University of Lavras, Campus Universitário, 3037, 37200-000, Lavras, MG, Brazil

✉ teo@dqf.ufla.br

☎ (+55) 35- 3829-1891

☎ (+55) 35- 3829-1271

1975, Castro and Figueroa-Villar 2002, Bartosova et al. 2005, Gonçalves et al. 2006, 2010, Ekström et al. 2007, Worek et al. 2007, Delfino and Figueroa-Villar 2009, Delfino et al. 2009, Ramalho et al. 2010).

In the present work, in order to contribute to a better understanding of the mechanism of action of oximes (Fig. 1), we proposed the structures and evaluated *in silico* the affinities of 10 new oximes, derived from pyridine-imidazol bicycled systems, for the HssAChE active site inhibited by the neurotoxic agent tabun. The softwares MVD[®] and Spartan[®] were used to estimate the values of the oximes' affinity (measured by the docking scores). The 10 oximes were studied together with the standard oximes pralidoxime (2-PAM), trimedoxime (TMB-4), and obidoxime as references. Additionally the reactivation constants of 2PAM and its *ortho*, *meta* and *para* isomers were calculated according to a procedure formerly established by Ramalho et. al. (2010). We evaluated the influence of the oxime group position as a substituent in the structures for 2-PAM and the 10 new oximes and, also, the influence of the oximes *syn-anti* isomery in the docking scores values for all the molecules studied. Our results suggest that the affinities of the 10 new oximes for the HssAChE active site are better than all 2PAM's isomers and some were quite similar to TMB-4's. It was also observed that *meta*-2-PAM could present better affinities for the HssAChE active site than *para* and *ortho*-2-PAM, and that 62.5% of the *anti* isomers presented slightly better affinities for the HssAChE active site when compared to the *syn* isomers.

MATERIALS AND METHODS

Docking energy calculations

The structure of the HssAChE used was that phosphorylated by tabun proposed and optimized by Gonçalves et al. (2006) and complexed with toxogonine, using as template the structure reported by Kryger et al. (2000) deposited in the Proteins Data Bank (Bernstein et al. 1977, Berman et al. 2000) under the PDB code IB41. The water molecules were removed using the program MVD[®], the three-dimensional structures of the oximes (Fig. 2) were built and optimized with the software Spartan Pro 5[®] and their partial atomic charges calculated by the PM3 semi-empirical method. The compounds were docked into the HssAChE binding site using MVD[®] (Thomsen et al. 2006) according to instructions which considered all the protein residues as flexible. Binding sites were restricted within spheres with radius of 8 Å, centered at the toxogonine

binding site in the protein complex and enclosing all the active site residues. Due to the stochastic nature of the ligand-protein docking search algorithm, about 10 runs were conducted and 30 docking solutions were retained for each ligand. The best superimposing poses related to toxogonine, were chosen to the analysis performed in this work.

DFT Studies

QM/MM techniques were performed to determine the preferred route for the reactivation process. On the technical side, we applied a procedure combining docking technique and DFT calculations at the QM/MM interface for the enzymatic mechanism. The QM calculations were carried out in the Spartan08 (Hehre et al. 1999) and Gaussian98 (Frisch et al. 2001) packages. The QM region, which consisted of residues, neighbouring peptide bonds, link atoms, crystallographic water molecules, ligand and inhibitor, was confined into a sphere with a radius of 15 Å, centered at each oxime.

The initial coordinates for the heavy atoms were taken from the HssAChE 3D structure proposed by Gonçalves et al. (2006). All the transition states, intermediates and precursors involved were calculated. Each conformer was fully optimized at the DFT level with B3LYP/6-31G (Ramalho and Taft 2005). Furthermore, after each optimization, a force constant calculation was made in order to verify whether the optimized structures were indeed local minima (no imaginary frequencies) or transition states (one imaginary frequency).

RESULTS AND DISCUSSION

Docking results

The cavity (Fig. 3) of the HssAChE active site was calculated by MVD[®] as having 949,696 Å³. The results of the docking studies of the oximes studied inside this cavity, allowed us to identify the relevant H-Bonds that occur between each oxime and the amino acid residues of the active site in order to obtain the conformations adopted with these molecules, compare them to the conformations of toxogonine and thus get subsidies for the investigations performed in this work.

The values of the docking energies obtained for the best poses of the *syn-anti* isomers of the 10 new oximes, the *ortho*, *meta* and *para* isomers of 2-PAM, TMB-4 and obidoxime are presented in Fig. 4. Table 1 reports the H-bond energy values obtained for each ligand in the HssAChE active site and, also, the amino acid residues involved in H-bonds with them.

Fig. 1. Inhibition, desinhibition and ageing of acetylcholinesterase. X is the leaving group.

Fig. 2. Structures of the oximes studied.

Fig. 3. Cavity of the active site of *HssAChE* complexed with obidoxime.

An analysis of Fig. 4 indicates that all the 10 new oximes isomers presented docking energies better than the 2PAM isomers but worse than TMB-4 and obidoxime isomers. This result reflects the smaller size of 2PAM and the new oximes, avoiding a full superposition to obidoxime and interactions with the totality of residues in the cavity (Figs 5 and 6). However, oximes 2, 3, 5, 6 and 8 presented at least one isomer with similar docking values to TMB-4 isomers.

H-bond energy values obtained for most of the new oximes isomers are better than for 2PAM and quite similar to the values obtained for TMB-4 and obidoxime isomers (see Table 1). All the aminoacid residues observed forming H-bonds with 2PAM were also observed for the new oximes. The new oximes also were able to form H-bonds with some additional

Fig. 4. Comparative docking energies of the *syn-anti* isomers of the new oximes, 2PAM isomers, HI-6, TMB-4 and obidoxime.

Fig. 5. Superposition of oxime 6 to obidoxime.

Fig. 6. Superposition of TMB-4 to obidoxime.

residues not observed for 2PAM (Tyr68, Asp70, Tyr120, Glu281 and Phe291). All the residues observed making H-bonds with the new oximes were observed in TMB-4 and obidoxime isomers which were, also, able to make additional H-bonds with the residues: Gly117, Gly118, Glu198, Ser199, Ala200 and His443 (see Table 1).

Concerning the position of the oxime group in 2PAM, the *anti* isomer of *meta*-2PAM presented a slightly better docking energy value than *para* and *ortho*-2PAM isomers but similar H-bond value to *ortho*-2PAM. However the best result among the 2PAM *syn* isomers was observed for *ortho*-2PAM.

The presence of these amino acids with 2PAM, 2PAM1 and 2PAM2 compounds in *anti* and *syn* conformations, suggests that they have a direct influence on the mechanism of reactivation of *HssAChE* inhibited by the organophosphorus agent Tabun.

Finally, analysis of the effect of the *syn-anti* isomery on the oximes' affinities for the *HssAChE* active site showed that for 10 of the 16 oximes studied (62.5%), the *anti* isomers presented better docking values than the *syn* isomers. For the H-bonds energy values the result pointed to 50%.

Reaction

The kinetics of the AChE reactivation process is believed to occur in two steps: (1) the association of the oxime to the inhibited AChE and (2) the reactivation of AChE by the leaving of the oxime complexed to the neurotoxic agent (Fig. 1). The process of kinetic reactivation of AChE could be illustrated by Equation 1:

Where EI is the organophosphate-inhibited enzyme, Ox is the reactivator (oxime), E is the

Table 1. H-bond energy values in kcal/mol obtained and main aminoacid residues interacting with the ligands.

Oxime	<i>Anti</i>	<i>Syn</i>	Oxime	<i>Anti</i>	<i>Syn</i>
1	-7.2209	-10.7584	8	-12.0939	-3.9584
	Phe291 (1)	Phe291 (1)		Trp282 (1)	Tyr 120 (1)
	Arg292 (2)	Arg292 (4)		Ser294 (1)	Ser294 (1)
	Ser289 (1)	Ser294 (1)		Arg292 (3)	Gly118 (1)
	Gly118 (1)	Gly118 (1)		Glu281 (1)	Tyr120 (1)
	Tyr120 (1)	Tyr120 (1)		Gly118 (1)	Tyr129 (1)
	Tyr129 (1)	Tyr129 (1)		Tyr120 (1)	Glu198 (1)
	Glu198 (1)	Glu198 (1)		Tyr129 (1)	
2			9	Glu198 (1)	
	-10.3404	-8.6853		-7.9259	0.0528
	Glu281 (1)	Trp282 (1)		Tyr68 (1)	Trp282 (2)
	Arg292 (3)	Tyr68 (1)		Ser294 (1)	Ser294 (1)
	Trp282 (1)	Arg292 (3)		Arg292 (2)	Glu281 (1)
	Ser294 (1)	Ser 294 (1)		Gly118 (1)	Arg292 (3)
	Gly118 (1)	Gly118 (1)		Tyr120 (1)	Gly118 (1)
	Tyr120 (1)	Tyr120 (1)		Tyr129 (1)	Tyr120 (1)
3			10	Glu198 (1)	Tyr129 (1)
	-7.7793	-7.7927			Glu198 (1)
	Trp282 (1)	Trp282 (1)		-9.7289	-10.7966
	Ser289 (1)	Ser289 (1)		Asp70 (1)	Arg292 (3)
	Arg292 (3)	Arg292 (3)		Tyr120 (1)	Ser294 (2)
	Ser294 (3)	Ser294 (3)		Ser294 (2)	Gly118 (1)
	Glu281 (1)	Glu281 (1)		Gly118 (1)	Tyr120 (1)
	Gly118 (1)	Gly118 (1)		Tyr129 (1)	Tyr129 (1)
4			2PAM	Glu198 (1)	Glu198 (1)
	-6.7650	-9.7872			
	Asp70 (1)	Tyr120 (2)		-5.9830	-5.4829
	Phe291 (1)	Ser294 (2)		Trp282 (1)	Trp282 (1)
	Arg292 (3)	Gly118 (1)		Arg292 (3)	Arg292 (3)
	Gly118 (1)	Tyr129 (1)		Gly118 (1)	Ser294 (1)
	Tyr120 (1)	Glu198 (1)		Tyr120 (1)	Gly118 (1)
	Tyr129 (1)			Tyr129 (1)	Tyr120 (1)
5			2PAM1	Glu198 (1)	Tyr129 (1)
	-7.4288	-1.7457			Glu198 (1)
	Asp70 (1)	Trp282 (1)		-1.6366	3.4180
	Trp282 (1)	Arg292 (3)		Trp282 (1)	Trp282 (1)
	Arg292 (3)	Gly118 (1)		Ser289 (1)	Arg292 (1)
	Gly118 (1)	Tyr120 (1)		Arg292 (3)	Gly118 (1)
	Tyr120 (1)	Tyr129 (1)		Gly118 (1)	Tyr120 (1)
	Tyr129 (1)	Glu198 (1)		Tyr120 (1)	Tyr129 (1)
	Glu198 (1)			Tyr129 (1)	Glu198 (1)
				Glu198 (1)	

(Continues)

(Continues)

Oxime	<i>Anti</i>	<i>Syn</i>	Oxime	<i>Anti</i>	<i>Syn</i>
6	−11.0145	−8.1143	2PAM2	−5.7507	−5.8037
	Asp70 (2)	Trp282 (1)		Trp282 (1)	Trp282 (2)
	Ser294 (3)	Ser294 (3)		Arg292 (3)	Arg292 (2)
	Gly118 (1)	Arg292 (3)		Gly118 (1)	Ser294 (1)
	Tyr120 (1)	Gly118 (1)		Tyr120 (1)	Gly118 (1)
	Tyr129 (1)	Tyr120 (1)		Tyr129 (1)	Tyr120 (1)
	Glu198 (1)	Tyr129 (1)		Glu198 (1)	Tyr129 (1)
				Glu198 (1)	
7	−9.0249	−7.3808			
	Trp282 (2)	Trp282 (1)			
	Ser294 (2)	Ser294 (2)			
	Arg292 (2)	Arg292 (3)			
	Gly118 (1)	Gly118 (1)			
	Tyr120 (1)	Tyr120 (1)			
	Tyr129 (1)	Tyr129 (1)			
	Glu198 (1)				

reactivated enzyme, EIOx is the intermediate complex and I-Ox is the product. K_R and k_R are the dissociation constants, which represent the affinity of oximes for tabun-inhibited AChE, and the rate constant for the decomposition of the stable enzyme-inhibitor-reactivator complex, respectively.

Nowadays, despite the recent efforts at both theoretical and experimental levels to elucidate the reaction mechanism involved in the reactivation process, some relevant facts, such as the influence of the oxime's isomery, still need to be clarified. Fig. 7 displays the proposed reaction mechanism considering 2PAM with *syn* and *anti* conformation.

2PAM was selected for the mechanism study due to experimental evidence, which suggest that this oxime could be used as a lead compound in order to propose structures of new oximes, such as 2PAM1 and 2PAM2.

The *syn* and *anti* conformations present the hydroxyl group close to the phosphate group of tabun, with distance values around 3.14Å and 3.24Å respectively (Fig. 8). For 2PAM1 and 2PAM2, similar distances were obtained for amino acid residues from the protein, favouring then, the reaction process.

Fig. 7. Scheme of the reactivation mechanism of the acetylcholinesterase enzyme.

Fig. 8. Compound 2PAM in the *anti* and *syn* conformations, respectively.

Fig. 9. Intermolecular steric effect in the compound *anti*-2PAM.

From our theoretical calculations of the catalytic mechanism, we obtained the relative activation energy of the three isomers 2PAM, 2PAM1 and 2PAM2, in different conformations, *syn* and *anti*. Theoretical data from Table 2 show that 2PAM and 2PAM2 in the *syn* conformation perform lower activation energy values than the compounds in *anti* conformations. This means that those conformations revealed a lower energetic barrier for the reaction pathway.

The lower thermodynamic stability of the 2PAM isomers with *anti* conformation, can be rationalized due to the lower number of hydrogen interactions with the residues close to the active site (Trp282, Arg292, Gly118, Tyr120, Tyr129, Glu198). Regarding the compounds in the *syn* conformation (Trp282, Arg292, Ser294, Gly118, Tyr120, Tyr129, Glu198), we can observe a lower stability inside the active site, suggesting a less stable transition state (higher activation energy).

Table 2. Relative activation energy values of the studied oximes.

Compounds	$\Delta\Delta E^\ddagger$ (kcal mol ⁻¹)
2PAM <i>syn</i>	31.98
2PAM <i>anti</i>	34.05
2PAM1 <i>syn</i>	10.30
2PAM1 <i>anti</i>	0.00
2PAM2 <i>syn</i>	23.88
2PAM2 <i>anti</i>	32.48

Turning now to the 2PAM1, our theoretical data point out that the *syn* conformation presents a steric hindrance between the aromatic hydrogen and the hydroxyl group in the same compound. The same scenario did not occur with 2PAM1 in the *anti* conformation, leaving the hydroxyl group free to interact with the phosphate group of the inhibitor. This molecule, then, possesses significant internal degrees of freedom, leading consequently to a lower activation energy value for the compound in the *anti* conformation.

Besides *anti* and *syn* conformations, the *ortho*, *meta* and *para* substituent orientations in the compounds, can also affect the transition state stability. In this way, we noticed that the compounds 2PAM and 2PAM2 have a methyl group in *ortho*-orientation, generating a higher steric hindrance with the hydroxyl group in the *anti* conformation, resulting in a higher activation energy value in relation to all other compounds, as reported in Fig. 9 and Table 2.

CONCLUSION

From the results obtained and discussed here it is possible to conclude that: 1) oximes derived from pyridine imidazol bicycled systems are worth synthesizing and testing *in vitro* as HssAChE reactivators and are expected to present similar experimental results as TMB-4; 2) the position of the oxime group as substituent, in the six or the five membered ring, on these molecules seems not to have a determinant influence on their affinities for the HssAChE active site; 3) *meta*-2PAM should be considered further as an HssAChE reactivator in experimental studies and 4) it is important, also, to consider a deeper investigation of the influence of the

anti isomers in experimental studies of oximes as HssAChE reactivators.

ACKNOWLEDGMENTS

The authors wish to thank the Brazilian financial agencies CNPq, FAPERJ, FAPEMIG and CAPES/PRODEFESA for financial support and the Military Institute of Engineering for the physical infrastructure and working space. This work was also supported by Ministry of Defense (Czech Republic) – FVZUO0000604 (KK).

REFERENCES

- Bartosova L, Kuca K, Jun D, Kunesova G. Bispyridinium oximes as antidotal treatment of cyclosarin poisoning – *in vitro* and *in vivo* testing. Internat J Tox. 24: 399–402, 2005.
- Bay E, Krop S, Yates LF. Chemotherapeutic effectiveness of 1,1'-trimethylene bis (4-formylpyridinium bromide) di-oxime (TMB-4) in experimental anticholinesterase poisoning. Proc Soc Exp Biol Med. 98: 107–110, 1958.
- Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat T N, Weissig H, Shindyalov IN, Bourne PE. The protein data bank. Nucleic Acids Res. 28: 235–242, 2000.
- Bernstein FC, Koetzle TF, Williams GJ, Meyer EE, Brice MD, Rodgers JR, Kennard O, Shimanouchi T, Tasumi M. The protein data bank. A computer-based archival file for macromolecular structures. J Mol Biol. 112: 535–542, 1977.
- Castro AT, Figueroa-Villar JD. Molecular structure, conformational analysis and charge distribution of

- pralidoxime: Ab initio and DFT studies. *Int J Quantum Chem.* 89: 135–146, 2002.
- Delfino RT, Figueroa-Villar JD. Nucleophilic reactivation of sarin-inhibited acetylcholinesterase: A molecular modeling study. *J Phys Chem B.* 113: 8402–8411, 2009.
- Delfino RT, Ribeiro TS, Figueroa-Villar JD. Organophosphorus compounds as chemical warfare agents: a review. *J Braz Chem Soc.* 20: 407–428, 2009.
- Ekström F J, Astot C, Pang YP. Novel nerve-agent antidote design based on crystallographic and mass spectrometric analyses of tabun-conjugated acetylcholinesterase in complex with antidotes, *Clin Pharmacol Ther.* 82: 282–293, 2007.
- Frisch MJ, Trucks GW, Schlegel HB, Scuseria ES, Pople JA. Gaussian 98 (Revision A.11). Gaussian Pittsburgh 2001.
- Gonçalves AS, França TCC, Wilter A, Figueroa-Villar JD. Molecular dynamics of the interaction of pralidoxime and deazapralidoxime with acetylcholinesterase inhibited by the neurotoxic agent tabun. *J Braz Chem Soc.* 17: 968–975, 2006.
- Gonçalves AS, França TCC, Figueroa-Villar JD, Pascutti PG. Conformational analysis of toxogonine, TMB-4 and HI-6 using PM6 and RM1 methods, *J Braz Chem Soc.* 21: 179–184, 2010.
- Hehre WJ, Deppmeier BJ, Klunzinger PE. PC SPARTAN Pro. Wavefunction Inc., Irvine 1999.
- Kassa J, Kuca K, Bartosova L, Kunesova G. The development of new structural analogues of oximes for the antidotal treatment of poisoning by nerve agents and the comparison of their reactivating and therapeutic efficacy with currently available oximes. *Curr Org Chem.* 11: 267–283, 2007.
- Kryger G, Harel M, Giles K, Toker L, Velan B, Lazar A, Kronman C, Barak D, Ariel N, Silman I, Shafferman A, Sussman JL. Structures of recombinant native and E202Q mutant human acetylcholinesterase complexed with the snake-venom toxin fasciculatin-II. *Acta Crystallogr.* 56: 1385–1394, 2000.
- Ramalho TC, Taft CA. Thermal and solvent effects on the NMR and UV parameters of some bioreductive drugs. *J Chem Physics.* 123: 054319–054328, 2005.
- Ramalho TC, França TCC, Rennó MN, Guimarães AP, da Cunha EFF, Kuča K. Development of new acetylcholinesterase reactivators: Molecular modeling versus *in vitro* data. *Chem Biol Inter.* 185: 73–77, 2010.
- Smirnova OI, E Gurina IL, Zhigalova V, Arestova LS. Toxicity and tolerance of toxogonin, *Farmakol Toksikol.* 38: 467–470, 1975.
- Thomsen R, Christensen MH. MolDock: A new technique for high-accuracy molecular docking. *J Med Chem.* 49: 3315–3321, 2006.
- Worek F, Aurbek N, Koller M, Becker C, Eyer P, Thiermann H. Kinetic analysis of reactivation and aging of human acetylcholinesterase inhibited by different phosphoramidates. *Biochem Pharmacol.* 73: 1807–1817, 2007.