

ORIGINAL ARTICLE

Reactivation study of pyridinium oximes for acetylcholinesterases inhibited by paraoxon or DFP

Tae-Hyuk Kim¹, Kyung-Ae Oh^{1,2}, No-Joong Park¹, No-Sang Park¹, Yeong Joon Kim², Eul Kyun Yum², Young-Sik Jung¹

¹ Medicinal Science Division, Korea Research Institute of Chemical Technology, P. O. Box 107, Yusong, Daejeon 305-606, Korea

² Department of Chemistry, Chungnam National University, Yusong, Daejeon 305-764, Korea

Received 30th October 2005.

Revised 14th December 2005.

Published online 20th February 2006.

Summary

We tested the potency to reactivate AChE inhibited by diisopropyl fluorophosphates (DFP) by using bis-pyridinium oxime reactivators connected with CH₂CH₂OCH₂CH₂ linkers between two pyridinium rings. The potency was strongly dependent on oxime functional groups, and the bis-oxime derivatives 1,1-[Oxybis(ethylene)]-bis[4-(hydroxyimino)methyl]pyridinium dibromide (1) and 1,1-[Oxybis(ethylene)]-bis[2-(hydroxyimino)methyl]pyridinium dichloride (2) are more potent than mono-oxime compounds 1-(4-hydroxyiminomethyl-1-pyridino)-5-(4-carbamoyl-1-pyridino)-3-oxapentane dibromide (3) and 1-(3-hydroxyiminomethyl-1-pyridino)-5-(4-carbamoyl-1-pyridino)-3-oxapentane dibromide (4). Not only is the number of oxime groups an important structural factor, but also their position. The *in vitro* reactivation ability of the most potent bis-pyridinium oxime 2 was further evaluated for the housefly (HF) AChE inhibited by DFP and the bovine red blood cell (RBC) AChE inhibited by paraoxon. The reactivation ability of oxime 2 at 5×10⁻³M concentration was almost 80% for HF-AChE inhibited by DFP and 82.1% for RBC-AChE inhibited by paraoxon.

Keywords: Paraoxon – DFP – Organophosphorus agents – Bis-pyridinium oxime reactivators – Acetylcholinesterase

INTRODUCTION

Organophosphorus nerve agents such as sarin, soman, and cyclosarin are extremely toxic chemi-


cals that were developed in secrecy for military use. It is well known that these organophosphorus agents exert their biological effects by inhibition of the enzyme acetylcholinesterase (AChE), which the active center serine hydroxyl group can attack the phosphorus atom of the organophosphorus agents to form a strong P-O bond (Scheme 1) (Marrs 1993). The inhibition of AChE increases the amount of acetylcholine (ACh) at central and peripheral sites of the nerve system. High doses of organophosphorus agents cause convulsions and paralysis of the respiratory muscles.

✉ Young-Sik Jung, Medicinal Science Division, Korea Research Institute of Chemical Technology, P. O. Box 107, Yusong, Daejeon 305-606, Korea

ysjung@kRICT.re.kr

+82-42-860-7135

+82-42-861-1291



Scheme 1. Inhibition of AChE by paraoxon and DFP

Several other organophosphorus agents such as parathion, malathion, and diazinon have also been developed (Figure 1), and became widely used as insecticides because of their low volatility and stability in aqueous solution (Taylor 1994). Among these organophosphorus insecticides, parathion probably has been responsible for more cases of accidental poisoning and death than any other organophosphorus insecticide (London and Myers 1995, Yacoub et. al. 1981).




Fig. 1. Structures of organophosphorus agents

After the organophosphorus compounds attach to AChE to inhibit it, AChE can be reactivated by oxime reactivators, and 2-PAM is the best well-known reactivator. After thorough study of many of the oxime reactivators, bis-pyridinium oximes such as TMB, Toxogonin, and HI-6 (Kassa et al. 1997, Kassa 1998, Kassa 2002) have been developed, and are used currently in many countries (Figure 2). In a previous paper (Kim et al. 2005), new bis-pyridinium oxime reactivators connected with a $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2$ linker between two pyridinium rings were designed and synthesized. During tests of their potency to reactivate AChE inhibited by cyclosarin, the bis-pyridinium oxime achieved

reactivation potency higher than 10% at the lower concentration 10^{-4}M .




Fig. 2. Currently used oxime reactivators

Although these compounds were not extraordinarily potent reactivators for AChE inhibited by cyclosarin, they could be effective in reactivation for AChE inhibited by other nerve agents or pesticides, because the reactivation potency of AChE reactivators depends on the organophosphorus agent used (Bajgar 2004, Dohnal et al. 2005, Kuca and Kassa 2003, Kuca et al. 2003).

In continuing our efforts to develop new oxime reactivators for AChE inhibited by organophosphorus agents, we are interested in the development of antidotes that have potent reactivation activity. Therefore, we tested the reactivation efficiency of the new oximes for

diisopropyl fluorophosphates (DFP)- or paraoxon-inhibited AChE (Figure 3).




Fig. 3. Pyridinium oxime compounds

MATERIAL AND METHODS

All new pyridinium oximes and HI-6 were prepared in our laboratory, and 2-PAM was purchased from Sigma-Aldrich. DFP and paraoxon are commercially available from Fluka and Sigma-Aldrich, respectively. Two kinds of AChE were used in this experiment; the first was extract from housefly (HF) head (Central Research Center, National Agricultural Cooperative Federation, Korea), and the other was bovine red blood cells (RBC) AChE, which were purchased from Sigma-Aldrich.

Determination of AChE activity

The enzyme activity was measured in a 96-well Microplate using a microplate reader (Benchmark Microplate Reader, BioRad) at 415 nm and 37 °C with acetylthiocholine (1 mM) as substrate and DTNB (1 mM) as chromogen in 0.05 M Tris-HCl buffer, pH 7.8 (Park and Kamble 2001) with a slight modification of Ellman's AChE assay method (Ellman et al. 1961). For RBC AChE, 1% of Triton X-100 (Sigma-Aldrich), we added a Tris-HCl buffer to preserve enzyme activity (Rosenberry and Scoggin 1984). The percentage reactivation of AChE activity was measured by the change of optical density per minute (OD/min) after correction for the control reaction.

AChE Inhibition and reactivation

AChE was inhibited with enough of the inhibitor to inactivate 99% for 10 minutes at room temperature. The concentration of DFP was 12.5 μM for HF AChE and 25 μM for RBC AChE, respectively, and 20 μM of paraoxon for both AChEs. To remove surplus inhibitor molecules after inhibition, the aqueous phase was separated by centrifugation after partitioning with two volumes of hexane (Worek et al. 1998). The collected solution containing phosphorylated AChE was incubated with various concentrations of 2-PAM or HI-6 for various reactivation times, respectively. Small molecules such as the reactivator and phosphorylated oxime were removed by filtration through a micro spin-column packed with Sephadex-G50 (Bio-Rad) (Luo et al. 1998). The AChE activity of the filtrate was measured in a 96-well microplate, and the percentage reactivation of AChE activity was calculated as previously described.

AChE Reactivation with newly synthesized oxime compounds

The reactivating capability of newly synthesized oxime compounds was evaluated against DFP or paraoxon-inhibited HF or RBC AChE, respectively, as previously described, and 5 mM of each oxime compound for 30 minutes for DFP-inhibited AChE and for 1 hour for paraoxon-inhibited AChE, respectively. The percentage reactivation of AChE was calculated as previously described.

RESULTS AND DISCUSSION

It is well known that 3 types of AChEs are generated after the post-transcriptional process of alternative splicing from the same origin (Taylor and Radic 1994). Each type of AChE exists in multiple forms, multimeric for nerve and muscle, dimeric for red blood cell, and monomeric for embryonic and tumor cells, respectively, and their structural difference appears only in the C-terminal extension with 40 residue peptides in contrast to the well-preserved functional subsites such as catalytic triad, acyl pocket, and hydrophobic subsite (Grisaru et al. 1999). House fly brain AChE and bovine red blood cells AChE were selected for this study as alternative forms of multimeric and dimeric AChEs. DFP has been primarily chosen for this study because of its close structural property to nerve gas (Taylor 2001). Parathion itself is inactive in inhibiting AChE *in vitro* in contrast to its metabolite paraoxon which is active. The sulfur-for-oxygen substitution is carried out predominantly in the liver by the mixed-function oxidases (Dauterman 1971, Butler and Murray

1997). This reaction is also carried out in insects, typically with more efficiency (O'Brien 1960, Oppenoorth 1972, Prestwich 1990). These two organophosphorus compounds have been used as representatives for the organophosphorus AChE inhibitor by many researchers (Gearhart et. al. 1994, Schwarz et. al. 1995, Krummer et. al. 2002,

Luo et. al. 2003). We have compared the reactivation potency of oximes (1-5) with two currently used AChE reactivators, 2-PAM and HI-6. As it is shown in Figure 4, the reactivation test was carried out for the DFP-inhibited housefly (HF).




Fig. 4. Reactivation potency of tested oximes for DFP-inhibited HF AChE

The potency was strongly dependent upon oxime functional groups, and bis-oxime derivatives (1, 2) are the most potent compared to mono-oxime compounds (3, 4). The position of the oxime group is also an important factor influencing the reactivation process, and compounds with the oxime group at position 2 on the pyridinium ring are the most potent reactivators for DFP-inhibited AChE. The potency of oxime 5 is same as that of oxime 1. From this preliminary result, bis-pyridinium oxime 2 is the most active among the prepared compounds and moreover oxime 2 is more potent than 2-PAM. Even though HI-6 is one of the most active reactivators against many organophosphorus nerve agents, it is no longer active against DFP at 5×10^{-3} M concentration. Therefore oxime 2 was selected for further evaluation of its reactivation activities. Figure 5 shows the reactivation potency of the oxime 2 for HF and RBC-AChE inhibited by DFP or paraoxon,

and compared with the potency of 2-PAM and HI-6. 2-PAM is quite potent for HF-AChE inhibited by DFP and RBC-AChE inhibited by paraoxon, whereas HI-6 shows very weak potency from all tests. Oxime 2 is more potent than 2-PAM, and is especially potent for HF-AChE inhibited by DFP and RBC-AChE inhibited by paraoxon.

In summary, in this *in vitro* reactivation evaluation of the oxime compounds, we found the bis-pyridinium oxime 2 is a strong reactivator for HF-AChE inhibited by DFP and RBC-AChE inhibited by paraoxon.

ACKNOWLEDGEMENT

This work was supported by grants from the Ministry of Commerce, Industry and Energy in Korea.




Fig. 5. Reactivation potency of oxime 2 for DFP- and paraoxon-inhibited AChE

REFERENCES

- Bajgar J.: Organophosphates/nerve agent poisoning: mechanism of action, diagnosis, prophylaxis, and treatment. *Adv. Clin. Chem.* 38:151–216, 2004.
- Butler A.M., Murray M.: Biotransformation of parathion in human liver: participation of CYP3A4 and its inactivation during microsomal parathion oxidation. *J. Pharmacol. Exp. Ther.* 280: 966–973, 1997.
- Dauterman W.C.: Biological and nonbiological modification of organophosphorus compounds. *WHO Bull.* 44:133–150, 1971.
- Dohnal V., Kuca K., Jun D.: Prediction of a new broad-spectrum reactivator capable of reactivating acetylcholinesterase inhibited by nerve agents. *J. Appl. Biomed.* 3:139–145, 2005.
- Ellman G. L., Courtney K. D., Andres Jr. V., Featherstone B.C.: A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* 7:88–95, 1961.
- Gearhart J., Jepson G.W., Clewell H.J. et al.: Physiologically based pharmacokinetic model for the inhibition of acetylcholinesterase by organophosphate esters. *Environ. Health Perspect.* 102 (Suppl. 11):51–60, 1994.
- Grisaru D., Sternfeld M., Eldor A. et al.: Structural roles of acetylcholinesterase variants in biology and pathology. *Eur. J. Biochem.* 264:672–686, 1999.
- Kassa J., Cabal J., Bajgar J., Szinicz L.: The choice: HI-6, pralidoxime or obidoxime against nerve agents? *ASA Newsl.* 97:16–18, 1997.
- Kassa J.: A comparison of the therapeutic efficacy of conventional and modern oximes against supralethal doses of highly toxic organophosphates in mice. *Acta Med. (Hradec Kralove)* 41: 19–21, 1998.
- Kassa J.: Review of oximes in the antidotal treatment of poisoning by organophosphorus nerve agents. *J. Toxicol. Clin. Toxicol.* 40:803–816, 2002.
- Kim T.H., Kuca K., Jun D., Jung Y.S.: Design and synthesis of new bis-pyridinium oximes as cyclosarin-inhibited acetylcholinesterase reactivators. *Bioorg. Med. Chem. Lett.* 15:2914–2917, 2005.
- Krummer S., Thiermann H., Worek F., Eyer P.: Equipotent cholinesterase reactivation in vitro by the nerve agent antidotes HI 6 dichloride and HI 6 dimethanesulfonate. *Arch. Toxicol.* 76:589–595, 2002.
- Kuca K., Kassa J.: A comparison of the ability of a new bispyridinium oxime-1-(4-hydroxyiminomethylpyridinium)-4-(4-carbamoylpyridinium) butane dibromide and currently used oximes to reactivate nerve agent-inhibited rat brain acetylcholinesterase by in vitro methods. *J. Enzyme Inhib. Med. Chem.* 18:529–535, 2003.
- Kuca K., Patocka J., Cabal J.: Reactivation of organophosphate inhibited acetylcholinesterase activity by -bis-(4-

- hydroxyiminomethylpyridinium) alkanes in vitro. *J. Appl. Biomed.* 1:207–211, 2003.
- London L., Myers J.E.: Agrichemical usage patterns and workplace exposure in the major farming sectors in the Southern Region of South Africa. *S. Afr. J. Sci.* 91:515–22, 1995.
- Luo C., Ashani Y., Doctor B.P.: Acceleration of oxime-induced reactivation of organophosphate-inhibited fetal bovine serum acetylcholinesterase by monoquaternary and bisquaternary ligands. *Mol. Pharmacol.* 53:718–726, 1998.
- Luo C., Leader H., Radic Z. et al.: Two possible orientations of the HI-6 molecule in the reactivation of organophosphate-inhibited acetylcholinesterase. *Biochem. Pharmacol.* 66:387–392, 2003.
- Marrs T.C.: Organophosphate poisoning. *Pharmacol. Ther.* 58: 51–66, 1993.
- O'Brien R.D.: Allelic genes in the house fly producing modified enzymes that cause organophosphorus resistance. *Science* 132:298–299, 1960.
- Oppenoorth F.J.: Degradation and activation of organophosphorus insecticides and resistance in insects. In Mushtaq A.K. (eds.): *Toxicology, Biodegradation and Efficacy of Livestock Pesticides*, W.O. Haufe Publ., Amsterdam 1972, pp. 73–92.
- Park N.J., Kamble S.T.: Decapitation impacting effect of topically applied chlorpyrifos on acetylcholinesterase and general esterases in susceptible and resistant German cockroaches (*Dictyoptera: Blattellidae*). *J. Econ. Entomol.* 94:499–505, 2001.
- Prestwich G.D.: Proinsecticides: metabolically activated toxicants. In Hodgson E., Kuhr R.J. (eds.): *Safer Insecticides, Development and Use*, Martin Dekker, New York 1990, pp. 281–335.
- Rosenberry T.L., Scoggin D.M.: Structure of human erythrocyte acetylcholinesterase. Characterization of intersubunit disulfide bonding and detergent interaction. *J. Biol. Chem.* 259:5643–5652, 1984.
- Schwarz M., Loewenstein-Lichtenstein Y., Glick D. et al.: Successive organophosphate inhibition and oxime reactivation reveals distinct responses of recombinant human cholinesterase variants. *Brain Res. Mol. Brain Res.* 31:101–110, 1995.
- Taylor P.: Anticholinergic agents. In: *The Pharmacological Basis of Therapeutics*. Hill, New York 2001, pp. 175–191.
- Taylor P., Radic Z.: The cholinesterases: from genes to proteins. *Annu. Rev. Pharmacol. Toxicol.* 34:281–320, 1994.
- Worek F., Widmann R., Knopff O., Szinicz L.: Reactivating potency of obidoxime, pralidoxime, HI-6 and Hlo-7 in human erythrocyte acetylcholinesterase inhibited by highly toxic organophosphorus compound. *Arch. Toxicol.* 72:237–243, 1998.
- Yacoub M., Skouri H., Amamou M., Besancenot F.: Intoxications aiguës par les insecticides organo-phosphoré. *J. Toxicol. Med.* 1:165–188, 1981.